


Convergence rates





Convergence rates



In order to compare perfomance of algorithms we need to define a terminology for different types of convergence.

Let $r_k = \{ \|x_k - x^*\|_2 \}$ be a sequence in \mathbb{R}^n that converges to zero.

♥ C 0 4

In order to compare perfomance of algorithms we need to define a terminology for different types of convergence. Let $r_k = \{\|x_k - x^*\|_2\}$ be a sequence in \mathbb{R}^n that converges to zero.

We can define the *linear* convergence in a two different forms:

$$||x_{k+1} - x^*||_2 \le Cq^k$$
 or $||x_{k+1} - x^*||_2 \le q||x_k - x^*||_2$,


♥ ი ⊘

In order to compare performance of algorithms we need to define a terminology for different types of convergence.

Let $r_k = \{||x_k - x^*||_2\}$ be a sequence in \mathbb{R}^n that converges to zero.

We can define the *linear* convergence in a two different forms:

on permand
$$\|x_{k+1}-x^*\|_2 \leq Cq^k$$



for all sufficiently large k. Here $q \in (0,1)$ and $0 < C < \infty$. This means that the distance to the solution x^* decreases at each iteration by at least a constant factor bounded away from 1. Note, that sometimes this type of convergence is also called *exponential* or *geometric*. The q is called the convergence rate.

In order to compare perfomance of algorithms we need to define a terminology for different types of convergence. Let $r_k = \{\|x_k - x^*\|_2\}$ be a sequence in \mathbb{R}^n that converges to zero.

We can define the *linear* convergence in a two different forms:

$$||x_{k+1} - x^*||_2 \le Cq^k$$
 or $||x_{k+1} - x^*||_2 \le q||x_k - x^*||_2$,

for all sufficiently large k. Here $q \in (0,1)$ and $0 < C < \infty$. This means that the distance to the solution x^* decreases at each iteration by at least a constant factor bounded away from 1. Note, that sometimes this type of convergence is also called *exponential* or *geometric*. The q is called the convergence rate.

i Question

Suppose, you have two sequences with linear convergence rates $q_1=0.1$ and $q_2=0.7$, which one is faster?

 $f \to \min_{x,y,z}$

⊕ ∩ ∅

i Example

Let us have the following sequence:

$$r_k = \frac{1}{2^k}$$

One can immediately conclude, that we have a linear convergence with parameters $q=\frac{1}{2}$ and C=1.

Convergence rates

i Example

Let us have the following sequence:

$$r_k = \frac{1}{2^k}$$

One can immediately conclude, that we have a linear convergence with parameters $q=\frac{1}{2}$ and C=1.

i Question

Determine the convergence of the following sequence

$$r_k = \frac{3}{2^k}$$

Sub and super

ogó runcinas

Sublinear convergence

If the sequence r_k converges to zero, but does not have linear convergence, the convergence is said to be sublinear. Sometimes we can consider the following class of sublinear convergence:

$$||x_{k+1} - x^*||_2 \le Ck^q,$$

where q < 0 and $0 < C < \infty$. Note, that sublinear convergence means, that the sequence is converging slower, than any geometric progression.

Superlinear convergence

cheoxnuteutias

The convergence is said to be *superlinear* if it converges to zero faster, than any linearly convergent sequence.

Sub and super

cydruneines, CX-T6

1 K!

REPLACHEL

Sublinear convergence

If the sequence r_k converges to zero, but does not have linear convergence, the convergence is said to be sublinear. Sometimes we can consider the following class of sublinear convergence:

$$||x_{k+1} - x^*||_2 < Ck^q$$

is converging slower, than

where q<0 and $0< C<\infty$. Note, that sublinear convergence means, that the sequence is converging slower, than any geometric progression.

Superlinear convergence

The convergence is said to be *superlinear* if it converges to zero faster, than any linearly convergent sequence.

 $f \to \min_{x,y}$

Convergence rates

i Theorem

Let $(r_k)_{k=m}^{\infty}$ be a sequence of non-negative numbers converging to zero, and let $\alpha := \limsup_{k \to \infty} r_k^{1/k}$. (Note that $\alpha > 0$.)

(a) If $0 \le \alpha < 1$, then $(r_k)_{k=m}^{\infty}$ converges linearly with constant α .

Proof.

i Theorem

Let $(r_k)_{k=m}^{\infty}$ be a sequence of non-negative numbers converging to zero, and let $\alpha := \limsup_{k \to \infty} r_k^{1/k}$. (Note that $\alpha > 0$.)

- (a) If $0 \le \alpha < 1$, then $(r_k)_{k=m}^{\infty}$ converges linearly with constant α .
- (b) In particular, if $\alpha = 0$, then $(r_k)_{k=m}^{\infty}$ converges superlinearly.

Proof.

i Theorem

Let $(r_k)_{k=m}^{\infty}$ be a sequence of non-negative numbers converging to zero, and let $\alpha:=\limsup_{k\to\infty}r_k^{1/k}$. (Note that $\alpha>0$.)

- (a) If $0 \le \alpha < 1$, then $(r_k)_{k=m}^{\infty}$ converges linearly with constant α .
- (b) In particular, if $\alpha = 0$, then $(r_k)_{k=m}^{\infty}$ converges superlinearly.
- (c) If $\alpha = 1$, then $(r_k)_{k=m}^{\infty}$ converges sublinearly.

Proof.

 $f \to \min_{x \in \mathbb{R}^n}$

i Theorem

Let $(r_k)_{k=m}^{\infty}$ be a sequence of non-negative numbers converging to zero, and let $\alpha:=\limsup_{k\to\infty}r_k^{1/k}$. (Note that $\alpha\geq 0$.)

- (a) If $0 \le \alpha < 1$, then $(r_k)_{k=m}^{\infty}$ converges linearly with constant α . (b) In particular, if $\alpha = 0$, then $(r_k)_{k=m}^{\infty}$ converges superlinearly.
- (c) If $\alpha = 1$, then $(r_k)_{k=m}^{\infty}$ converges sublinearly.
- (d) The case $\alpha > 1$ is impossible.

Proof.

Convergence rates

♥ C €

i Theorem

Let $(r_k)_{k=m}^{\infty}$ be a sequence of non-negative numbers converging to zero, and let $\alpha := \limsup_{k \to \infty} r_k^{1/k}$. (Note that $\alpha > 0$.)

- (a) If $0 \le \alpha < 1$, then $(r_k)_{k=m}^{\infty}$ converges linearly with constant α .
- (b) In particular, if $\alpha = 0$, then $(r_k)_{k=m}^{\infty}$ converges superlinearly.
- (c) If $\alpha = 1$, then $(r_k)_{k=m}^{\infty}$ converges sublinearly. (d) The case $\alpha > 1$ is impossible.

Proof.

1. let us show that if $(r_k)_{k=m}^{\infty}$ converges linearly with constant $0 \le \beta < 1$, then necessarily $\alpha \le \beta$.

Indeed, by the definition of the constant of linear convergence, for any $\varepsilon > 0$ satisfying $\beta + \varepsilon < 1$, there exists C>0 such that $r_k\leq C(\beta+\varepsilon)^k$ for all $k\geq m$.

From this, $r_k^{1/k} \leq C^{1/k}(\beta + \varepsilon)$ for all $k \geq m$. Passing to the limit as $k \to \infty$ and using $C^{1/k} \to 1$, we obtain $\alpha \leq \beta + \varepsilon$. Given the arbitrariness of ε , it follows that $\alpha \leq \beta$.

i Theorem

(d) The case $\alpha > 1$ is impossible.

Let $(r_k)_{k=m}^\infty$ be a sequence of non-negative numbers converging to zero, and let $\alpha:=\limsup_{k\to\infty}r_k^{1/k}$. (Note that $\alpha\geq 0$.)

(a) If $0\leq \alpha<1$, then $(r_k)_{k=m}^\infty$ converges linearly with constant α .

(b) In particular, if $\alpha=0$, then $(r_k)_{k=m}^\infty$ converges superlinearly.

(c) If $\alpha=1$, then $(r_k)_{k=m}^\infty$ converges sublinearly.

Proof.

1. let us show that if $(r_k)_{k=m}^{\infty}$ converges linearly with constant $0 \le \beta < 1$, then necessarily $\alpha \le \beta$.

Indeed, by the definition of the constant of linear convergence, for any $\varepsilon>0$ satisfying $\beta+\varepsilon<1$, there exists C>0 such that $r_k\leq C(\beta+\varepsilon)^k$ for all $k\geq m$.

From this, $r_k^{1/k} \leq C^{1/k}(\beta + \varepsilon)$ for all $k \geq m$. Passing to the limit as $k \to \infty$ and using $C^{1/k} \to 1$, we obtain $\alpha \leq \beta + \varepsilon$. Given the arbitrariness of ε , it follows that $\alpha \leq \beta$.

2. Thus, in the case $\alpha=1$, the sequence $(r_k)_{k=m}^{\infty}$ cannot have linear convergence according to the above result (proven by contradiction). Since, nevertheless, $(r_k)_{k=m}^{\infty}$ converges to zero, it must converge sublinearly.



3. Now consider the case $0 \le \alpha < 1$. Let $\varepsilon > 0$ be an arbitrary number such that $\underline{\alpha + \varepsilon < 1}$.

3. Now consider the case $0 \le \alpha < 1$. Let $\varepsilon > 0$ be an arbitrary number such that $\alpha + \varepsilon < 1$.

Npulle:
$$r_{x} = \frac{1}{K^{x}}$$

$$d = \lim_{K \to \infty} \sup_{K \to \infty} \left(\frac{1}{K^{x}} \right) = \lim_{K \to \infty} \sup_{K \to \infty} \frac{1}{K^{x}}$$

 $f \to \min_{x,y,z}$

Convergence rates

⊕ ი ⊘

3. Now consider the case $0 \le \alpha < 1$. Let $\varepsilon > 0$ be an arbitrary number such that $\alpha + \varepsilon < 1$.

According to the properties of the limsup, there exists $N \geq m$ such that $r_k^{1/k} \leq \alpha + \varepsilon$ for all $k \geq N$.

Convergence rates

3. Now consider the case $0 \le \alpha < 1$. Let $\varepsilon > 0$ be an arbitrary number such that $\alpha + \varepsilon < 1$.

According to the properties of the limsup, there exists $N \geq m$ such that $r_k^{1/k} \leq \alpha + \varepsilon$ for all $k \geq N$. Hence, $r_k \leq (\alpha + \varepsilon)^k$ for all $k \geq N$. Therefore, $(r_k)_{k=m}^\infty$ converges linearly with parameter $\alpha + \varepsilon$ (it does not

matter that the inequality is only valid from the number N).


d+E + unerce

3. Now consider the case $0 \le \alpha < 1$. Let $\varepsilon > 0$ be an arbitrary number such that $\alpha + \varepsilon < 1$.

According to the properties of the limsup, there exists $N \ge m$ such that $r_k^{1/k} \le \alpha + \varepsilon$ for all $k \ge N$.

Hence, $r_k \leq (\alpha + \varepsilon)^k$ for all $k \geq N$. Therefore, $(r_k)_{k=m}^{\infty}$ converges linearly with parameter $\alpha + \varepsilon$ (it does not matter that the inequality is only valid from the number N).

Due to the arbitrariness of ε , this means that the constant of linear convergence of $(r_k)_{k=m}^{\infty}$ does not exceed α .



BEd



d < B

Convergence rates

3. Now consider the case $0 \le \alpha < 1$. Let $\varepsilon > 0$ be an arbitrary number such that $\alpha + \varepsilon < 1$.

According to the properties of the limsup, there exists $N \ge m$ such that $r_k^{1/k} \le \alpha + \varepsilon$ for all $k \ge N$.

Hence, $r_k \leq (\alpha + \varepsilon)^k$ for all $k \geq N$. Therefore, $(r_k)_{k=m}^{\infty}$ converges linearly with parameter $\alpha + \varepsilon$ (it does not matter that the inequality is only valid from the number N).

Due to the arbitrariness of ε , this means that the constant of linear convergence of $(r_k)_{k=m}^{\infty}$ does not exceed α .

Since, as shown above, the constant of linear convergence cannot be less than α , this means that the constant of linear convergence of $(r_k)_{k=m}^{\infty}$ is exactly α .

4. Finally, let's show that the case $\alpha > 1$ is impossible.



3. Now consider the case $0 \le \alpha < 1$. Let $\varepsilon > 0$ be an arbitrary number such that $\alpha + \varepsilon < 1$.

According to the properties of the limsup, there exists $N \ge m$ such that $r_k^{1/k} \le \alpha + \varepsilon$ for all $k \ge N$.

Hence, $r_k \leq (\alpha + \varepsilon)^k$ for all $k \geq N$. Therefore, $(r_k)_{k=m}^{\infty}$ converges linearly with parameter $\alpha + \varepsilon$ (it does not matter that the inequality is only valid from the number N).

Due to the arbitrariness of ε , this means that the constant of linear convergence of $(r_k)_{k=m}^{\infty}$ does not exceed α .

Since, as shown above, the constant of linear convergence cannot be less than α , this means that the constant of linear convergence of $(r_k)_{k=m}^{\infty}$ is exactly α .

4. Finally, let's show that the case $\alpha > 1$ is impossible.



3. Now consider the case $0 \le \alpha < 1$. Let $\varepsilon > 0$ be an arbitrary number such that $\alpha + \varepsilon < 1$.

According to the properties of the limsup, there exists $N \ge m$ such that $r_k^{1/k} \le \alpha + \varepsilon$ for all $k \ge N$.

Hence, $r_k \leq (\alpha + \varepsilon)^k$ for all $k \geq N$. Therefore, $(r_k)_{k=m}^{\infty}$ converges linearly with parameter $\alpha + \varepsilon$ (it does not matter that the inequality is only valid from the number N).

Due to the arbitrariness of ε , this means that the constant of linear convergence of $(r_k)_{k=m}^{\infty}$ does not exceed α .

Since, as shown above, the constant of linear convergence cannot be less than α , this means that the constant of linear convergence of $(r_k)_{k=m}^{\infty}$ is exactly α .

4. Finally, let's show that the case $\alpha > 1$ is impossible.

Indeed, suppose $\alpha > 1$. Then from the definition of limsup, it follows that for any $N \ge m$, there exists $k \ge N$ such that $r_k^{1/k} > 1$, and, in particular, $r_k > 1$.

3. Now consider the case $0 \le \alpha < 1$. Let $\varepsilon > 0$ be an arbitrary number such that $\alpha + \varepsilon < 1$.

According to the properties of the limsup, there exists $N \ge m$ such that $r_k^{1/k} \le \alpha + \varepsilon$ for all $k \ge N$.

Hence, $r_k \leq (\alpha + \varepsilon)^k$ for all $k \geq N$. Therefore, $(r_k)_{k=m}^{\infty}$ converges linearly with parameter $\alpha + \varepsilon$ (it does not matter that the inequality is only valid from the number N).

Due to the arbitrariness of ε , this means that the constant of linear convergence of $(r_k)_{k=m}^{\infty}$ does not exceed α .

Since, as shown above, the constant of linear convergence cannot be less than α , this means that the constant of linear convergence of $(r_k)_{k=m}^{\infty}$ is exactly α .

4. Finally, let's show that the case $\alpha > 1$ is impossible.

Indeed, suppose $\alpha > 1$. Then from the definition of limsup, it follows that for any $N \ge m$, there exists $k \ge N$ such that $r_k^{1/k} > 1$, and, in particular, $r_k > 1$.

But this means that r_k has a subsequence that is bounded away from zero. Hence, $(r_k)_{k=m}^{\infty}$ cannot converge to zero, which contradicts the condition.

 $f \to \min_{x,y,z}$ Convergence rates

Let $\{r_k\}_{k=m}^{\infty}$ be a sequence of strictly positive numbers converging to zero. Let

$$q = \lim_{k \to \infty} \frac{r_{k+1}}{r_k}$$

• If there exists q and $0 \le q < 1$, then $\{r_k\}_{k=m}^{\infty}$ has linear convergence with constant q.

Let $\{r_k\}_{k=m}^{\infty}$ be a sequence of strictly positive numbers converging to zero. Let

$$q = \lim_{k \to \infty} \frac{r_{k+1}}{r_k}$$

- If there exists q and $0 \le q < 1$, then $\{r_k\}_{k=m}^{\infty}$ has linear convergence with constant q.
- In particular, if q=0, then $\{r_k\}_{k=m}^{\infty}$ has superlinear convergence.



Let $\{r_k\}_{k=m}^{\infty}$ be a sequence of strictly positive numbers converging to zero. Let

$$q = \lim_{k \to \infty} \frac{r_{k+1}}{r_k}$$

- If there exists q and $0 \le q < 1$, then $\{r_k\}_{k=m}^{\infty}$ has linear convergence with constant q.
- In particular, if q=0, then $\{r_k\}_{k=m}^{\infty}$ has superlinear convergence.
- If q does not exist, but $q=\lim_{k\to\infty}\sup_k\frac{r_{k+1}}{r_k}<1$, then $\{r_k\}_{k=m}^\infty$ has linear convergence with a constant not exceeding q.



Let $\{r_k\}_{k=m}^{\infty}$ be a sequence of strictly positive numbers converging to zero. Let

$$q = \lim_{k \to \infty} \frac{r_{k+1}}{r_k}$$

- If there exists q and $0 \le q < 1$, then $\{r_k\}_{k=m}^{\infty}$ has linear convergence with constant q.
- In particular, if q=0, then $\{r_k\}_{k=m}^{\infty}$ has superlinear convergence.
- If q does not exist, but $q = \lim_{k \to \infty} \sup_k \frac{r_{k+1}}{r_k} < 1$, then $\{r_k\}_{k=m}^{\infty}$ has linear convergence with a constant not exceeding q.
- If $\lim_{k\to\infty}\inf_k\frac{r_{k+1}}{r_k}=1$, then $\{r_k\}_{k=m}^\infty$ has sublinear convergence.

Let $\{r_k\}_{k=m}^{\infty}$ be a sequence of strictly positive numbers converging to zero. Let

$$q = \lim_{k \to \infty} \frac{r_{k+1}}{r_k}$$

- If there exists q and $0 \le q < 1$, then $\{r_k\}_{k=m}^{\infty}$ has linear convergence with constant q.
- In particular, if q=0, then $\{r_k\}_{k=m}^{\infty}$ has superlinear convergence.
- If q does not exist, but $q=\lim_{k\to\infty}\sup_k\frac{r_{k+1}}{r_k}<1$, then $\{r_k\}_{k=m}^\infty$ has linear convergence with a constant not exceeding q.
- If $\lim_{k\to\infty}\inf_k\frac{r_{k+1}}{r_k}=1$, then $\{r_k\}_{k=m}^\infty$ has sublinear convergence.
- The case $\lim_{k\to\infty}\inf_k\frac{r_{k+1}}{r_{k}}>1$ is impossible.



Let $\{r_k\}_{k=m}^{\infty}$ be a sequence of strictly positive numbers converging to zero. Let

$$q = \lim_{k \to \infty} \frac{r_{k+1}}{r_k}$$

- If there exists q and $0 \le q < 1$, then $\{r_k\}_{k=m}^{\infty}$ has linear convergence with constant q.
- In particular, if q=0, then $\{r_k\}_{k=m}^{\infty}$ has superlinear convergence.
- If q does not exist, but $q=\lim_{k\to\infty}\sup_k\frac{r_{k+1}}{r_k}<1$, then $\{r_k\}_{k=m}^\infty$ has linear convergence with a constant not exceeding q.
- If $\lim_{k\to\infty}\inf_k\frac{r_{k+1}}{r_k}=1$, then $\{r_k\}_{k=m}^\infty$ has sublinear convergence.
- The case $\lim_{k \to \infty} \inf_k \frac{r_{k+1}}{r_k} > 1$ is impossible.
- In all other cases (i.e., when $\lim_{k\to\infty}\inf_k\frac{r_{k+1}}{r_k}<1\leq\lim_{k\to\infty}\sup_k\frac{r_{k+1}}{r_k}$) we cannot claim anything concrete about the convergence rate $\{r_k\}_{k=m}^\infty$.



Ratio test lemma

i Theorem

Let $(r_k)_{k=m}^{\infty}$ be a sequence of strictly positive numbers. (The strict positivity is necessary to ensure that the ratios $\frac{r_{k+1}}{r_k}$, which appear below, are well-defined.) Then

$$\liminf_{k\to\infty}\frac{r_{k+1}}{r_k}\le \liminf_{k\to\infty}r_k^{1/k}\le \limsup_{k\to\infty}r_k^{1/k}\le \limsup_{k\to\infty}\frac{r_{k+1}}{r_k}.$$

Ratio test lemma

i Theorem

Let $(r_k)_{k=m}^{\infty}$ be a sequence of strictly positive numbers. (The strict positivity is necessary to ensure that the ratios $\frac{r_{k+1}}{r_{\nu}}$, which appear below, are well-defined.) Then

$$\liminf_{k\to\infty}\frac{r_{k+1}}{r_k}\leq \liminf_{k\to\infty}r_k^{1/k}\leq \limsup_{k\to\infty}r_k^{1/k}\leq \limsup_{k\to\infty}\frac{r_{k+1}}{r_k}.$$

Proof.

1. The middle inequality follows from the fact that the liminf of any sequence is always less than or equal to its limsup. Let's prove the last inequality; the first one is proved analogously.

⊕ O

Ratio test lemma

i Theorem

Let $(r_k)_{k=m}^{\infty}$ be a sequence of strictly positive numbers. (The strict positivity is necessary to ensure that the ratios $\frac{r_{k+1}}{r_k}$, which appear below, are well-defined.) Then

$$\liminf_{k\to\infty}\frac{r_{k+1}}{r_k}\leq \liminf_{k\to\infty}r_k^{1/k}\leq \limsup_{k\to\infty}r_k^{1/k}\leq \limsup_{k\to\infty}\frac{r_{k+1}}{r_k}.$$

Proof.

- 1. The middle inequality follows from the fact that the liminf of any sequence is always less than or equal to its limsup. Let's prove the last inequality; the first one is proved analogously.
- 2. Denote $L:=\limsup_{k\to\infty} \frac{r_{k+1}}{r_k}$. If $L=+\infty$, then the inequality is obviously true, so let's assume L is finite.

Note that $L\geq 0$, since the ratio $\frac{r_{k+1}}{r_k}$ is positive for all $k\geq m$. Let $\varepsilon>0$ be an arbitrary number. According to the properties of limsup, there exists $N \geq m$ such that $\frac{r_{k+1}}{r_i} \leq L + \varepsilon$ for all $k \geq N$. From here,

 $r_{k+1} \leq (L+\varepsilon)r_k$ for all $k \geq N$. Applying induction, we get $r_k \leq (L+\varepsilon)^{k-N}r_N$ for all $k \geq N$. Let $C:=(L+\varepsilon)^{-N}r_N$. Then $r_k\leq C(L+\varepsilon)^k$ for all $k\geq N$, from which $r_k^{1/k}\leq C^{1/k}(L+\varepsilon)$. Taking the limsup as $k \to \infty$ and using $C^{1/k} \to 1$, we get $\limsup_{k \to \infty} r_k^{1/k} \le L + \varepsilon$. Given the arbitrariness of ε , it follows that $\limsup_{k\to\infty} r_k^{1/k} < L.$

 $J \to \min_{x,y,z}$ Line search

0

Suppose, we have a problem of minimization of a function $f(x): \mathbb{R} \to \mathbb{R}$ of scalar variable:

$$f(x)\to \min_{x\in\mathbb{R}}$$


Metogles oghowlphoro noucks

⊕ ი

Suppose, we have a problem of minimization of a function $f(x): \mathbb{R} \to \mathbb{R}$ of scalar variable:

$$f(x) \to \min_{x \in \mathbb{R}}$$

Sometimes, we refer to the similar problem of finding minimum on the line segment [a,b]:



Suppose, we have a problem of minimization of a function $f(x): \mathbb{R} \to \mathbb{R}$ of scalar variable:

$$f(x) \to \min_{x \in \mathbb{R}}$$

Sometimes, we refer to the similar problem of finding minimum on the line segment [a, b]:

$$f(x) \to \min_{x \in [a,b]} \qquad f(x) = f(x) : \mathbb{R} \to \mathbb{R}$$

i Example

Typical example of line search problem is selecting appropriate stepsize for gradient descent algorithm:

$$x_{k+1} = x_k - \alpha \nabla f(x_k)$$

 $\alpha = \operatorname{argmin} f(x_{k+1})$

Suppose, we have a problem of minimization of a function $f(x): \mathbb{R} \to \mathbb{R}$ of scalar variable:

$$f(x) \to \min_{x \in \mathbb{R}}$$

Sometimes, we refer to the similar problem of finding minimum on the line segment $\left[a,b\right]$:

$$f(x) \to \min_{x \in [a,b]}$$

i Example

Typical example of line search problem is selecting appropriate stepsize for gradient descent algorithm:

$$x_{k+1} = x_k - \alpha \nabla f(x_k)$$
$$\alpha = \operatorname{argmin} f(x_{k+1})$$

The line search is a fundamental optimization problem that plays a crucial role in solving complex tasks. To simplify the problem, let's assume that the function, f(x), is unimodal, meaning it has a single peak or valley.

Unimodal function

i Definition

Function f(x) is called **unimodal** on [a,b], if there is $x_* \in [a,b]$, that $f(x_1) > f(x_2) \quad \forall a \leq x_1 < x_2 < x_*$ and $f(x_1) < f(x_2) \quad \forall x_* < x_1 < x_2 \leq b$



⇔ റ ഉ

Unimodal function

i Definition

Function f(x) is called **unimodal** on [a,b], if there is $x_* \in [a,b]$, that $f(x_1) > f(x_2) \quad \forall a \leq x_1 < x_2 < x_*$ and $f(x_1) < f(x_2) \quad \forall x_* < x_1 < x_2 \leq b$





Figure 2: Examples of unimodal functions

Let f(x) be unimodal function on [a,b]. Than if $x_1 < x_2 \in [a,b]$, then:

• if $f(x_1) \le f(x_2) \to x_* \in [a, x_2]$



Let f(x) be unimodal function on [a,b]. Than if $x_1 < x_2 \in [a,b]$, then:

- if $f(x_1) \le f(x_2) \to x_* \in [a, x_2]$
- if $f(x_1) \ge f(x_2) \to x_* \in [x_1, b]$

Let f(x) be unimodal function on [a,b]. Than if $x_1 < x_2 \in [a,b]$, then:

- if $f(x_1) \le f(x_2) \to x_* \in [a, x_2]$
- if $f(x_1) \ge f(x_2) \to x_* \in [x_1, b]$

Let f(x) be unimodal function on [a, b]. Than if $x_1 < x_2 \in [a, b]$, then:


- if $f(x_1) < f(x_2) \to x_* \in [a, x_2]$
- if $f(x_1) > f(x_2) \to x_* \in [x_1, b]$

Proof Let's prove the first statement. On the contrary, suppose that $f(x_1) \leq f(x_2)$, but $x^* > x_2$. Then necessarily $x_1 < x_2 < x^*$ and by the unimodality of the function f(x) the inequality: $f(x_1) > f(x_2)$ must be satisfied. We have obtained a contradiction.

Let f(x) be unimodal function on [a, b]. Than if $x_1 < x_2 \in [a, b]$, then:

- if $f(x_1) \le f(x_2) \to x_* \in [a, x_2]$
- if $f(x_1) > f(x_2) \to x_* \in [x_1, b]$


Proof Let's prove the first statement. On the contrary, suppose that $f(x_1) \leq f(x_2)$, but $x^* > x_2$. Then necessarily $x_1 < x_2 < x^*$ and by the unimodality of the function f(x) the inequality: $f(x_1) > f(x_2)$ must be satisfied. We have obtained a contradiction.



Let f(x) be unimodal function on [a, b]. Than if $x_1 < x_2 \in [a, b]$, then:

- if $f(x_1) < f(x_2) \to x_* \in [a, x_2]$
- if $f(x_1) > f(x_2) \to x_* \in [x_1, b]$


Proof Let's prove the first statement. On the contrary, suppose that $f(x_1) \leq f(x_2)$, but $x^* > x_2$. Then necessarily $x_1 < x_2 < x^*$ and by the unimodality of the function f(x) the inequality: $f(x_1) > f(x_2)$ must be satisfied. We have obtained a contradiction.



Let f(x) be unimodal function on [a,b]. Than if $x_1 < x_2 \in [a,b]$, then:

- if $f(x_1) \le f(x_2) \to x_* \in [a, x_2]$ • if $f(x_1) \ge f(x_2) \to x_* \in [x_1, b]$
- if $f(x_1) \ge f(x_2) \to x_* \in [x_1, t_1]$

Proof Let's prove the first statement. On the contrary, suppose that $f(x_1) \le f(x_2)$, but $x^* > x_2$. Then necessarily $x_1 < x_2 < x^*$ and by the unimodality of the function f(x) the inequality: $f(x_1) > f(x_2)$ must be satisfied. We have obtained a contradiction.



We aim to solve the following problem:

$$f(x) \to \min_{x \in [a,b]}$$

We divide a segment into two equal parts and choose the one that contains the solution of the problem using the values of functions, based on the key property described above. Our goal after one iteration of the method is to halve the solution region.




Figure 3: Dichotomy method for unimodal function



We measure the function value at the middle of the line segment





Figure 4: Dichotomy method for unimodal function

In order to apply the key property we perform another measurement.






Figure 5: Dichotomy method for unimodal function



We select the target line segment. And in this case we are lucky since we already halved the solution region. But that is not always the case.




Figure 6: Dichotomy method for unimodal function

Dichotomy methodLet's consider another unimodal function.




Figure 7: Dichotomy method for unimodal function

Dichotomy methodMeasure the middle of the line segment.




Figure 8: Dichotomy method for unimodal function

Dichotomy method Get another measurement.




Figure 9: Dichotomy method for unimodal function

Select the target line segment. You can clearly see, that the obtained line segment is not the half of the initial one. It is $\frac{3}{4}(b-a)$. So to fix it we need another step of the algorithm.




Figure 10: Dichotomy method for unimodal function

After another additional measurement, we will surely get

$$\frac{2}{3}\frac{3}{4}(b-a) = \frac{1}{2}(b-a)$$




Figure 11: Dichotomy method for unimodal function



To sum it up, each subsequent iteration will require at most two function value measurements.




Figure 12: Dichotomy method for unimodal function

Dichotomy method. Algorithm

```
def binary_search(f, a, b, epsilon):
    c = (a + b) / 2
    while abs(b - a) > epsilon:
        y = (a + c) / 2.0
        if f(y) \leq f(c):
            b = c
            c = y
        else:
            z = (b + c) / 2.0
            if f(c) \ll f(z):
                a = y
                b = z
            else:
                a = c
                c = z
    return c
```

The length of the line segment on k+1-th iteration:

$$\Delta_{k+1} = b_{k+1} - a_{k+1} = \frac{1}{2^k} (b - a)$$



େ ଚେଡ

The length of the line segment on k + 1-th iteration:

$$\Delta_{k+1} = b_{k+1} - a_{k+1} = \frac{1}{2^k}(b-a)$$

For unimodal functions, this holds if we select the middle of a segment as an output of the iteration x_{k+1} :

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{1}{2^{k+1}}(b-a) \leq (0.5)^{k+1} \cdot (b-a)$$

$$|x_{k+1} - x_*| \leq \frac{\Delta_{k+1}}{2} \leq \frac{\Delta_{k+$$

The length of the line segment on k + 1-th iteration:

$$\Delta_{k+1} = b_{k+1} - a_{k+1} = \frac{1}{2^k}(b-a)$$

For unimodal functions, this holds if we select the middle of a segment as an output of the iteration x_{k+1} :

$$|x_{k+1} - x_*| \le \frac{\Delta_{k+1}}{2} \le \frac{1}{2^{k+1}} (b-a) \le (0.5)^{k+1} \cdot (b-a)$$

Note, that at each iteration we ask oracle no more, than 2 times, so the number of function evaluations is $N=2\cdot k$, which implies:

$$|x_{k+1} - x_*| \le (0.5)^{\frac{N}{2} + 1} \cdot (b - a) \le (0.707)^N \frac{b - a}{2}$$

The length of the line segment on k + 1-th iteration:

$$\Delta_{k+1} = b_{k+1} - a_{k+1} = \frac{1}{2^k}(b-a)$$

For unimodal functions, this holds if we select the middle of a segment as an output of the iteration x_{k+1} :

$$|x_{k+1} - x_*| \le \frac{\Delta_{k+1}}{2} \le \frac{1}{2^{k+1}} (b-a) \le (0.5)^{k+1} \cdot (b-a)$$

Note, that at each iteration we ask oracle no more, than 2 times, so the number of function evaluations is $N=2\cdot k$, which implies:

$$|x_{k+1} - x_*| \le (0.5)^{\frac{N}{2} + 1} \cdot (b - a) \le (0.707)^N \frac{b - a}{2}$$

By marking the right side of the last inequality for ε , we get the number of method iterations needed to achieve ε accuracy:

$$K = \left\lceil \log_2 \frac{b - a}{\varepsilon} - 1 \right\rceil$$

 $f \to \min_{x,y,z}$ Line search

Golden selection

The idea is quite similar to the dichotomy method. There are two golden points on the line segment (left and right) and the insightful idea is, that on the next iteration one of the points will remain the golden point.




Figure 13: Key idea, that allows us to decrease function evaluations



Golden section. Algorithm

```
def golden_search(f, a, b, epsilon):
    tau = (sqrt(5) + 1) / 2
    y = a + (b - a) / tau**2
    z = a + (b - a) / tau
    while b - a > epsilon:
        if f(y) \leq f(z):
            b = z
            z = y
            v = a + (b - a) / tau**2
        else:
            a = v
            y = z
            z = a + (b - a) / tau
    return (a + b) / 2
```

Golden section. Bounds

$$|x_{k+1} - x_*| \le b_{k+1} - a_{k+1} = \left(\frac{1}{\tau}\right)^{N-1} (b-a) \approx 0.618^k (b-a),$$

where $au = \frac{\sqrt{5}+1}{2}$.

ullet The geometric progression constant **more** than the dichotomy method - 0.618 worse than 0.5





Golden section. Bounds

$$|x_{k+1} - x_*| \le b_{k+1} - a_{k+1} = \left(\frac{1}{\tau}\right)^{N-1} (b-a) \approx 0.618^k (b-a),$$

where $\tau = \frac{\sqrt{5}+1}{2}$.

- ullet The geometric progression constant more than the dichotomy method 0.618 worse than 0.5
- ullet The number of function calls **is less** than for the dichotomy method 0.707 worse than 0.618 (for each iteration of the dichotomy method, except for the first one, the function is calculated no more than 2 times, and for the gold method - no more than one)





Successive parabolic interpolation

Sampling 3 points of a function determines unique parabola. Using this information we will go directly to its minimum. Suppose, we have 3 points $x_1 < x_2 < x_3$ such that line segment $[x_1, x_3]$ contains minimum of a function f(x). Then, we need to solve the following system of equations:



Successive parabolic interpolation

Sampling 3 points of a function determines unique parabola. Using this information we will go directly to its minimum. Suppose, we have 3 points $x_1 < x_2 < x_3$ such that line segment $[x_1, x_3]$ contains minimum of a function f(x). Then, we need to solve the following system of equations:

$$ax_i^2 + bx_i + c = f_i = f(x_i), i = 1, 2, 3$$

Note, that this system is linear, since we need to solve it on a,b,c. Minimum of this parabola will be calculated as:

Successive parabolic interpolation

Sampling 3 points of a function determines unique parabola. Using this information we will go directly to its minimum. Suppose, we have 3 points $x_1 < x_2 < x_3$ such that line segment $[x_1, x_3]$ contains minimum of a function f(x). Then, we need to solve the following system of equations:

$$ax_i^2 + bx_i + c = f_i = f(x_i), i = 1, 2, 3$$

Note, that this system is linear, since we need to solve it on a,b,c. Minimum of this parabola will be calculated as:

$$u = -\frac{b}{2a} = x_2 - \frac{(x_2 - x_1)^2 (f_2 - f_3) - (x_2 - x_3)^2 (f_2 - f_1)}{2[(x_2 - x_1)(f_2 - f_3) - (x_2 - x_3)(f_2 - f_1)]}$$

Note, that if $f_2 < f_1, f_2 < f_3$, than u will lie in $[x_1, x_3]$


 $ightarrow \min_{x,y,z}$ Line search

Successive parabolic interpolation. Algorithm ¹

```
def parabola_search(f, x1, x2, x3, epsilon):
   f1, f2, f3 = f(x1), f(x2), f(x3)
   while x3 - x1 > epsilon:
       u = x^2 - ((x^2 - x^1) * * 2 * (f^2 - f^3) - (x^2 - x^3) * * 2 * (f^2 - f^1)) / (2 * ((x^2 - x^1) * (f^2 - f^3) - (x^2 - x^3) * (f^2 - f^1)))
       fn = f(n)
                                                          SCIPY. Minimize_scolor
       if x2 <= u:
          if f2 <= fu:
              x1. x2. x3 = x1. x2. u
              f1, f2, f3 = f1, f2, fu
           else:
              x1. x2. x3 = x2. u. x3
              f1. f2, f3 = f2, fu, f3
       else:
           if fu <= f2:
              x1, x2, x3 = x1, u, x2
                                                                Kondunagua 301. (1).
napadon. uhtepn.
              f1. f2. f3 = f1. fu. f2
           else.
              x1, x2, x3 = u, x2, x3
             f1, f2, f3 = fu, f2, f3
   return (x1 + x3) / 2
```

 $f \to \min_{x,y,z}$ Line search

 $^{^{1}}$ The convergence of this method is superlinear, but local, which means, that you can take profit from using this method only near some neighbour of optimum. Here is the proof of superlinear convergence of order 1.32.



Inexact line search

Sometimes it is enough to find a solution, which will approximately solve out problem. This is very typical scenario for mentioned stepsize selection problem

$$x_{k+1} = x_k - \alpha \nabla f(x_k)$$

$$\alpha = \operatorname{argmin} f(x_{k+1})$$

$$= \int \left(\chi_{k} - \chi \nabla f(x_k) - \chi_{k+1} \right)$$

େ ଚେ ଚ

Inexact line search

Sometimes it is enough to find a solution, which will approximately solve out problem. This is very typical scenario for mentioned stepsize selection problem

$$x_{k+1} = x_k - \alpha \nabla f(x_k)$$
$$\alpha = \operatorname{argmin} f(x_{k+1})$$

Consider a scalar function $\phi(\alpha)$ at a point x_k :

consider a scalar function
$$\phi(\alpha)$$
 at a point x_k .
$$\phi(\alpha) = f(x_k - \alpha \nabla f(x_k)) |\alpha| \ge 0$$

$$\phi(\alpha) = f(x_k - \alpha \nabla f(x_k)), \alpha \ge 0$$

$$\phi(\alpha) = f(x_k - \alpha \nabla f(x_k)), \alpha \ge 0$$

$$(d) = \psi(0) - \psi(0)$$

$$\varphi(0) + \frac{\partial \varphi}{\partial \lambda}(0) \cdot (\lambda - 0)$$

Inexact line search

Sometimes it is enough to find a solution, which will approximately solve out problem. This is very typical scenario for mentioned stepsize selection problem

$$x_{k+1} = x_k - \alpha \nabla f(x_k)$$
$$\alpha = \operatorname{argmin} f(x_{k+1})$$

Consider a scalar function $\phi(\alpha)$ at a point x_k :

$$\phi(\alpha) = f(x_k - \alpha \nabla f(x_k)), \alpha \ge 0$$

The first-order approximation of $\phi(\alpha)$ near $\alpha=0$ is:

$$\phi(\alpha) \approx f(x_k) - \alpha \nabla f(x_k)^T \nabla f(x_k)$$




Figure 14: Illustration of Taylor approximation of $\phi_0^I(\alpha)$

Inexact line search. Sufficient Decrease

The inexact line search condition, known as the Armijo condition, states that α should provide sufficient decrease in the function f, satisfying:

$$f(x_k - \alpha \nabla f(x_k)) \le f(x_k) - c_1 \cdot \alpha \nabla f(x_k)^T \nabla f(x_k)$$

Inexact line search. Sufficient Decrease

The inexact line search condition, known as the Armijo condition, states that α should provide sufficient decrease in the function f, satisfying:

$$f(x_k - \alpha \nabla f(x_k)) \le f(x_k) - c_1 \cdot \alpha \nabla f(x_k)^T \nabla f(x_k)$$

for some constant $c_1 \in (0,1)$. Note that setting $c_1=1$ corresponds to the first-order Taylor approximation of $\phi(\alpha)$. However, this condition can accept very small values of α , potentially slowing down the solution process. Typically, $c_1 \approx 10^{-4}$ is used in practice.



Inexact line search. Sufficient Decrease

The inexact line search condition, known as the Armijo condition, states that α should provide sufficient decrease

in the function f, satisfying: $f(x_k - \alpha \nabla f(x_k)) \le f(x_k) - c_1 \cdot \alpha \nabla f(x_k)^T \nabla f(x_k)$

$$f(x_k - \alpha \nabla f(x_k)) \le f(x_k) - c_1 \cdot \alpha \nabla f(x_k)^T \nabla f(x_k)$$

$$c \text{ some constant } c_1 \in (0, 1) \text{ Note that setting } c_1 = 1$$

for some constant $c_1 \in (0,1)$. Note that setting $c_1 = 1$ corresponds to the first-order Taylor approximation of $\phi(\alpha)$. However, this condition can accept very small values of α , potentially slowing down the solution process.

Typically, $c_1 \approx 10^{-4}$ is used in practice.

i Example

problem, choosing an appropriate c_1 value is crucial. For instance, in a machine learning model training scenario, an improper c_1 might lead to either very slow convergence or missing the minimum.

If f(x) represents a cost function in an optimization




Figure 15: Illustration of sufficient decrease condition with coefficient c_1

Inexact line search. Goldstein Conditions

Consider two linear scalar functions $\phi_1(\alpha)$ and $\phi_2(\alpha)$:

$$\phi_1(\alpha) = f(x_k) - c_1 \alpha \|\nabla f(x_k)\|^2$$

$$\phi_2(\alpha) = f(x_k) - c_2 \alpha \|\nabla f(x_k)\|^2$$


Inexact line search. Goldstein Conditions

Consider two linear scalar functions $\phi_1(\alpha)$ and $\phi_2(\alpha)$:

$$\phi_1(\alpha) = f(x_k) - c_1 \alpha \|\nabla f(x_k)\|^2$$

$$\phi_2(\alpha) = f(x_k) - c_2 \alpha \|\nabla f(x_k)\|^2$$

The Goldstein-Armijo conditions locate the function $\phi(\alpha)$ between $\phi_1(\alpha)$ and $\phi_2(\alpha)$. Typically, $c_1 = \rho$ and $c_2 = 1 - \rho$, with $\rho \in (0, 0.5)$.






Figure 16: Illustration of Goldstein conditions

Inexact line search. Curvature Condition

To avoid excessively short steps, we introduce a second criterion:

$$-\nabla f(x_k - \alpha \nabla f(x_k))^T \nabla f(x_k) \ge c_2 \nabla f(x_k)^T (-\nabla f(x_k))$$

Inexact line search. Curvature Condition

To avoid excessively short steps, we introduce a second criterion:

$$-\nabla f(x_k - \alpha \nabla f(x_k))^T \nabla f(x_k) \ge c_2 \nabla f(x_k)^T (-\nabla f(x_k))$$

for some $c_2 \in (c_1, 1)$. Here, c_1 is from the Armijo condition. The left-hand side is the derivative $\nabla_{\alpha} \phi(\alpha)$, ensuring that

conditions form the Wolfe conditions.

the slope of $\phi(\alpha)$ at the target point is at least c_2 times the initial slope $\nabla_{\alpha}\phi(\alpha)(0)$. Commonly, $c_2\approx 0.9$ is used for Newton or quasi-Newton methods. Together, the sufficient decrease and curvature




Figure 17: Illustration of curvature condition

⊕ ೧

Inexact line search. Wolfe Condition

$$-\nabla f(x_k - \alpha \nabla f(x_k))^T \nabla f(x_k) \ge c_2 \nabla f(x_k)^T (-\nabla f(x_k))$$

Together, the sufficient decrease and curvature conditions form the Wolfe conditions.

i Theorem

Let $f: \mathbb{R}^n \to \mathbb{R}$ be continuously differentiable, and let $\phi(\alpha) = f(x_k - \alpha \nabla f(x_k))$. Assume $\nabla f(x_k)^T p_k < 0$, where $p_k = -\nabla f(x_k)$, making p_k a descent direction. Also, assume f is bounded below along the ray $\{x_k + \alpha p_k \mid \alpha > 0\}$. We aim to show that for $0 < c_1 < c_2 < 1$, there exist intervals of step lengths satisfying the Wolfe conditions.




Figure 18: Illustration of Wolfe condition

Inexact line search. Wolfe Condition. Proof

1. Since $\phi(\alpha) = f(x_k + \alpha p_k)$ is bounded below and $l(\alpha) = f(x_k) + \alpha c_1 \nabla f(x_k)^T p_k$ is unbounded below (as $\nabla f(x_k)^T p_k < 0$), the graph of $l(\alpha)$ must intersect the graph of $\phi(\alpha)$ at least once. Let $\alpha' > 0$ be the smallest such value satisfying:

$$f(x_k + \alpha' p_k) \le f(x_k) + \alpha' c_1 \nabla f(x_k)^T p_k.$$
 (1)

This ensures the sufficient decrease condition is satisfied.

Line search

Inexact line search. Wolfe Condition. Proof

1. Since $\phi(\alpha) = f(x_k + \alpha p_k)$ is bounded below and $l(\alpha) = f(x_k) + \alpha c_1 \nabla f(x_k)^T p_k$ is unbounded below (as $\nabla f(x_k)^T p_k < 0$), the graph of $l(\alpha)$ must intersect the graph of $\phi(\alpha)$ at least once. Let $\alpha' > 0$ be the smallest such value satisfying:

$$f(x_k + \alpha' p_k) \le f(x_k) + \alpha' c_1 \nabla f(x_k)^T p_k.$$
 (1)

This ensures the **sufficient decrease condition** is satisfied.

2. By the Mean Value Theorem, there exists $\alpha'' \in (0, \alpha')$ such that:

$$f(x_k + \alpha' p_k) - f(x_k) = \alpha' \nabla f(x_k + \alpha'' p_k)^T p_k$$
. (2)

Substituting $f(x_k + \alpha' p_k)$ from (1) into (2), we have:

$$\alpha' \nabla f(x_k + \alpha'' p_k)^T p_k < \alpha' c_1 \nabla f(x_k)^T p_k.$$

Dividing through by $\alpha' > 0$, this simplifies to:

$$\nabla f(x_k + \alpha'' p_k)^T p_k \le c_1 \nabla f(x_k)^T p_k. \tag{3}$$



Inexact line search. Wolfe Condition. Proof

1. Since $\phi(\alpha) = f(x_k + \alpha p_k)$ is bounded below and

 $l(\alpha) = f(x_k) + \alpha c_1 \nabla f(x_k)^T p_k$ is unbounded below

(as $\nabla f(x_k)^T p_k < 0$), the graph of $l(\alpha)$ must intersect the graph of $\phi(\alpha)$ at least once. Let $\alpha' > 0$ be the smallest such value satisfying:

$$f(x_k + \alpha' p_k) \le f(x_k) + \alpha' c_1 \nabla f(x_k)^T p_k.$$
 (1)

This ensures the sufficient decrease condition is

satisfied. 2. By the Mean Value Theorem, there exists

$$f(x_k + \alpha' p_k) - f(x_k) = \alpha' \nabla f(x_k + \alpha'' p_k)^T p_k.$$
 (2)

 $\alpha'' \in (0, \alpha')$ such that:

Substituting $f(x_k + \alpha' p_k)$ from (1) into (2), we have:

$$\alpha' \nabla f(x_k + \alpha'' p_k)^T p_k \le \alpha' c_1 \nabla f(x_k)^T p_k.$$

Dividing through by $\alpha' > 0$, this simplifies to:

Dividing through by
$$\alpha > 0$$
, this simplifies to:
$$\nabla f(x_k + \alpha'' n_k)^T n_k < c_1 \nabla f(x_k)^T n_k.$$

(3)

3. Since $c_1 < c_2$ and $\nabla f(x_k)^T p_k < 0$, the inequality $c_1 \nabla f(x_k)^T p_k < c_2 \nabla f(x_k)^T p_k$ holds. This implies there exists α'' such that:

$$\nabla f(x_k + \alpha'' p_k)^T p_k \le c_2 \nabla f(x_k)^T p_k. \tag{Inequalities (3) and (4) together ensure the Wolfe}$$

conditions are satisfied.

Inexact line search. Wolfe Condition. Proof 1. Since $\phi(\alpha) = f(x_k + \alpha p_k)$ is bounded below and

 $l(\alpha) = f(x_k) + \alpha c_1 \nabla f(x_k)^T p_k$ is unbounded below

(as
$$\nabla f(x_k)^T p_k < 0$$
), the graph of $l(\alpha)$ must intersect the graph of $\phi(\alpha)$ at least once. Let $\alpha' > 0$ be the smallest such value satisfying:

$$f(x_k + \alpha' p_k) \le f(x_k) + \alpha' c_1 \nabla f(x_k)^T p_k.$$
 (1

This ensures the sufficient decrease condition is satisfied.

2. By the Mean Value Theorem, there exists

By the Mean Value Theorem, there exist
$$\alpha'' \in (0, \alpha')$$
 such that:

 $f(x_k + \alpha' p_k) - f(x_k) = \alpha' \nabla f(x_k + \alpha'' p_k)^T p_k$ (2) Substituting $f(x_k + \alpha' p_k)$ from (1) into (2), we have:

$$\alpha' \nabla f(x_k + \alpha'' p_k)^T p_k \le \alpha' c_1 \nabla f(x_k)^T p_k.$$

Dividing through by $\alpha' > 0$, this simplifies to:

$$\nabla f(x_k + \alpha'' p_k)^T p_k \le c_1 \nabla f(x_k)^T p_k.$$
 (3)

3. Since $c_1 < c_2$ and $\nabla f(x_k)^T p_k < 0$, the inequality $c_1 \nabla f(x_k)^T p_k < c_2 \nabla f(x_k)^T p_k$ holds. This implies there exists α'' such that:

$$\nabla f(x_k + \alpha'' p_k)^T p_k \le c_2 \nabla f(x_k)^T p_k.$$

Inequalities (3) and (4) together ensure the Wolfe conditions are satisfied. 4. For the strong Wolfe conditions, the curvature

condition:
$$\left|\nabla f(x_k + \alpha p_k)^T p_k\right| \le c_2 \left|\nabla f(x_k)^T p_k\right| \tag{5}$$

is met because $\nabla f(x_k + \alpha p_k)^T p_k$ is negative and bounded below by $c_2 \nabla f(x_k)^T p_k$.

(4)

Inexact line search. Wolfe Condition. Proof 1. Since $\phi(\alpha) = f(x_k + \alpha p_k)$ is bounded below and

 $l(\alpha) = f(x_k) + \alpha c_1 \nabla f(x_k)^T p_k$ is unbounded below

(as $\nabla f(x_k)^T p_k < 0$), the graph of $l(\alpha)$ must intersect the graph of $\phi(\alpha)$ at least once. Let $\alpha' > 0$ be the smallest such value satisfying:

$$f(x_k + \alpha' p_k) \le f(x_k) + \alpha' c_1 \nabla f(x_k)^T p_k.$$
 (1)

This ensures the sufficient decrease condition is satisfied.

2. By the Mean Value Theorem, there exists

$$\alpha'' \in (0, \alpha')$$
 such that:

$$f(x_k + \alpha' p_k) - f(x_k) = \alpha' \nabla f(x_k + \alpha'' p_k)^T p_k. \tag{2}$$

Substituting $f(x_k + \alpha' p_k)$ from (1) into (2), we have:

$$\alpha' \nabla f(x_k + \alpha'' p_k)^T p_k \le \alpha' c_1 \nabla f(x_k)^T p_k.$$
Dividing through by $\alpha' > 0$, this simplifies to:

$$\nabla f(x_k + \alpha'' p_k)^T p_k \le c_1 \nabla f(x_k)^T p_k.$$
 (3)

condition:

complete.

3. Since $c_1 < c_2$ and $\nabla f(x_k)^T p_k < 0$, the inequality $c_1 \nabla f(x_k)^T p_k < c_2 \nabla f(x_k)^T p_k$ holds. This implies there exists α'' such that:

$$\nabla f(x_k + \alpha'' p_k)^T p_k \le c_2 \nabla f(x_k)^T p_k.$$

Inequalities (3) and (4) together ensure the Wolfe conditions are satisfied. 4. For the strong Wolfe conditions, the curvature

$$\left| \nabla f(x_k + \alpha p_k)^T p_k \right| \le c_2 \left| \nabla f(x_k)^T p_k \right|$$
 (5) is met because $\nabla f(x_k + \alpha p_k)^T p_k$ is negative and

bounded below by $c_2 \nabla f(x_k)^T p_k$. Due to the smoothness of f, there exists an interval around α'' where the Wolfe conditions (and thus the strong Wolfe conditions) hold. Hence, the proof is

(4)

Backtracking line search is a technique to find a step size that satisfies the Armijo condition, Goldstein conditions, or other criteria of inexact line search. It begins with a relatively large step size and iteratively scales it down until a condition is met.

♥

Backtracking line search is a technique to find a step size that satisfies the Armijo condition, Goldstein conditions, or other criteria of inexact line search. It begins with a relatively large step size and iteratively scales it down until a condition is met.

Algorithm:

1. Choose an initial step size, α_0 , and parameters $\beta \in (0,1)$ and $c_1 \in (0,1)$.

Line search

♥

Backtracking line search is a technique to find a step size that satisfies the Armijo condition, Goldstein conditions, or other criteria of inexact line search. It begins with a relatively large step size and iteratively scales it down until a condition is met.

Algorithm:

- 1. Choose an initial step size, α_0 , and parameters $\beta \in (0,1)$ and $c_1 \in (0,1)$.
- 2. Check if the chosen step size satisfies the chosen condition (e.g., Armijo condition).

Line search

Backtracking line search is a technique to find a step size that satisfies the Armijo condition, Goldstein conditions, or other criteria of inexact line search. It begins with a relatively large step size and iteratively scales it down until a condition is met.

Algorithm:

- 1. Choose an initial step size, α_0 , and parameters $\beta \in (0,1)$ and $c_1 \in (0,1)$.
- 2. Check if the chosen step size satisfies the chosen condition (e.g., Armijo condition).
- 3. If the condition is satisfied, stop; else, set $\alpha := \beta \alpha$ and repeat step 2.

Backtracking line search is a technique to find a step size that satisfies the Armijo condition, Goldstein conditions, or other criteria of inexact line search. It begins with a relatively large step size and iteratively scales it down until a condition is met.

Algorithm:

- 1. Choose an initial step size, α_0 , and parameters $\beta \in (0,1)$ and $c_1 \in (0,1)$.
- 2. Check if the chosen step size satisfies the chosen condition (e.g., Armijo condition).
- 3. If the condition is satisfied, stop; else, set $\alpha := \beta \alpha$ and repeat step 2.

Backtracking line search is a technique to find a step size that satisfies the Armijo condition, Goldstein conditions, or other criteria of inexact line search. It begins with a relatively large step size and iteratively scales it down until a condition is met.

Algorithm:

- 1. Choose an initial step size, α_0 , and parameters $\beta \in (0,1)$ and $c_1 \in (0,1)$.
- 2. Check if the chosen step size satisfies the chosen condition (e.g., Armijo condition).

3. If the condition is satisfied, stop; else, set $\alpha := \beta \alpha$ and repeat step 2. The step size α is updated as

$$\alpha_{k+1} := \beta \alpha_k$$

in each iteration until the chosen condition is satisfied.

i Example

In machine learning model training, the backtracking line search can be used to adjust the learning rate. If the loss doesn't decrease sufficiently, the learning rate is reduced multiplicatively until the Armijo condition is met.



Numerical illustration




Figure 19: Comparison of different line search algorithms

Open In Colab 🖺



